

Online-only Data Supplement

Bleeding	Target INR	INR day 0	Procedure
MB	≤1.4	1.6	Nephrectomy
MB	≤1.4	1.7	Total hip arthroplasty
MB	≤1.4	1.3	Hemorrhoidectomy
CRNMB	≤1.4	1.6	Revision of lumbar arthrodesis
CRNMB	≤3.0	2.4	Excision of pilonidal cyst

Table S1. Postprocedural bleeding. MB: Major bleeding. CRNMB: Clinically relevant non-major bleeding.

Target INR: ≤1.4						
INR Day 0	Reached at Day 0	Reached at Day 1	Reached at Day 2	Reached at Day 3	Reached at Day 4	Reached at Day 5
1.5-1.9	0/14 = 0%	0/14 = 0%	8/14 = 57%	14/14 = 100%	14/14 = 100%	14/14 = 100%
2.0-2.4	0/23 = 0%	0/23 = 0%	11/23 = 48%	22/23 = 96%	23/23 = 100%	23/23 = 100%
2.5-2.9	0/25 = 0%	0/25 = 0%	9/25 = 36%	20/25 = 80%	23/25 = 92%	25/25 = 100%
3.0-3.4	0/14 = 0%	0/14 = 0%	2/14 = 14%	11/14 = 79%	13/14 = 93%	14/14 = 100%
3.5-3.9	0/2 = 0%	0/2 = 0%	0/2 = 0%	1/2 = 50%	2/2 = 100%	2/2 = 100%

Table S2. Number and percentage of patients that reached the target INR ≤1.4 day 0-5 after discontinuation of warfarin.

Target INR: ≤2.5						
INR Day 0	Reached at Day 0	Reached at Day 1	Reached at Day 2	Reached at Day 3	Reached at Day 4	Reached at Day 5
1.5-1.9	7/7 = 100%	7/7 = 100%	7/7 = 100%	7/7 = 100%	7/7 = 100%	7/7 = 100%
2.0-2.4	21/21 = 100%	21/21 = 100%	21/21 = 100%	21/21 = 100%	21/21 = 100%	21/21 = 100%
2.5-2.9	4/21 = 19%	18/21 = 86%	20/21 = 95%	21/21 = 100%	21/21 = 100%	21/21 = 100%
3.0-3.4	0/1 = 0%	1/1 = 100%	1/1 = 100%	1/1 = 100%	1/1 = 100%	1/1 = 100%
3.5-3.9	0/1 = 0%	0/1 = 0%	1/1 = 100%	1/1 = 100%	1/1 = 100%	1/1 = 100%

Table S3. Number and percentage of patients that reached the target INR ≤2.5 day 0-5 after discontinuation of warfarin.




Figure S1.

Influence of age on the maintenance dose of warfarin in patients with therapeutic interval 2.0-3.0.