

SUPPLEMENTARY MATERIAL FOR “Epidemiology of erysipelas and necrotizing soft tissue infections”

TABLE S1 Chief complaints and vital signs at arrival in adult acute non-trauma visits with erysipelas or necrotizing soft tissue infections in emergency/acute departments, Region of Southern Denmark, 2016–2018 stratified by age groups

	Total	18-69 years	≥70 years
Total, n (%)	2183 (100)	1105 (50.6)	1078 (49.4)
Sex			
Female	867 (39.7)	379 (34.3)	488 (45.3)
Male	1316 (60.3)	726 (65.7)	590 (54.7)
Symptoms			
Abdominal pain	34 (1.6)	24 (2.2)	10 (0.9)
Chest pain	17 (0.8)	7 (0.6)	10 (0.9)
Dyspnea	104 (4.8)	36 (3.3)	68 (6.3)
Fever	1102 (50.5)	602 (54.5)	500 (46.4)
Missing registration	78 (3.6)	44 (4.0)	34 (3.2)
Neurologic complaints	19 (0.9)	7 (0.6)	12 (1.1)
Others	135 (6.2)	56 (5.1)	79 (7.3)
Unspecific symptoms	694 (31.8)	329 (29.8)	365 (33.9)
Vital signs			
Systolic blood pressure, mean (SD), mmHg	136 (22.6)	135 (22.0)	137 (23.2)
Diastolic blood pressure, mean (SD), mmHg	74 (16.2)	78 (15.4)	71 (16.4)
Heart rate, mean (SD) bpm	90 (19.4)	93 (18.8)	88 (19.5)
Respiratory rate, mean (SD), brpm	19 (4.8)	19 (4.4)	20 (5.1)
Oxygen saturation, median (IQR), %	97 (95-98)	97 (96-99)	96 (94-98)
Glasgow coma scale, median (IQR)	15 (15-15)	15 (15-15)	15 (15-15)
Temperature, mean (SD), C	37.8 (1.1)	37.9 (1.1)	37.7 (1.1)

Data are numbers (%) unless otherwise noted.

bpm = beats per minute; brpm = breaths per minute; IQR = interquartile range; SD = standard deviation.