


## Supplementary material

Relates to: Retrospective analysis of patients referred to a single university hospital with Peyronie's disease.

**Supplementary Figure 1** - Flow chart over management strategy



i: the patient moved to Copenhagen, thereby changing the region.

ii: due to COVID

**Supplementary material**

Relates to: Retrospective analysis of patients referred to a single university hospital with Peyronie's disease.

**Supplementary Table 1 - Risk factors**

Parameter	All n = 26	No complication n = 21	Complicatio n n = 5
<b>Risk Factors</b>			
Alcohol use	24 (92.3)	20 (76.9)	3 (11.5)
<14 units/ week, n (%)	1 (3.8)	0	1 (3.8)
>14 units/ week, n (%)	1 (3.8)	1 (3.8)	1 (3.8)
No data, n (%)			
<b>Smoking</b>			
No smoking, n (%)	10 (38.3)	8 (30.7)	2 (7.6)
<10 cigarettes/ day, n (%)	3 (11.5)	3 (11.5)	0
10-20 cigarettes/ day, n (%)	3 (11.5)	2 (7.6)	1 (3.8)
20-30 cigarettes/ day, n (%)	0	0	0
40=< cigarettes/ day, n (%)	0	0	0
Former smoker, n (%), n (%)	9 (34.6)	8 (30.7)	1 (3.8)
No data, n (%)	1 (3.8)	0	1 (3.8)
<b>Diabetes Mellitus</b>			
Type I , n (%)	2 (7.6)	2 (7.6)	0
Type II, n (%)	2 (7.6)	2 (7.6)	0
None, n (%)*	22 (84.6)	17 (65.4)	5 (19.2)
No Data, n (%)	0	0	0