

26. Chelly JE, Greger J, Gebhard R et al. Continuous femoral blocks improve recovery and outcome of patients undergoing total knee arthroplasty. *J Arthroplasty* 2001;16:436-45.
27. Davies AF, Segar EP, Murdoch J et al. Epidural infusion or combined femoral and sciatic nerve blocks as perioperative analgesia for knee arthroplasty. *Br J Anaesth* 2004;93:368-74.
28. Morin AM, Kratz CD, Eberhart LH et al. Postoperative analgesia and functional recovery after total-knee replacement: comparison of a continuous posterior lumbar plexus (psoas compartment) block, a continuous femoral nerve block, and the combination of a continuous femoral and sciatic nerve block. *Reg Anesth Pain Med* 2005;30:434-45.
29. Pham DC, Gautheron E, Guilley J et al. The value of adding sciatic block to continuous femoral block for analgesia after total knee replacement. *Reg Anesth Pain Med* 2005;30:128-33.
30. Allen HW, Liu SS, Ware PD et al. Peripheral nerve blocks improve analgesia after total knee replacement surgery. *Anesth Analg* 1998;87:93-7.
31. Eledjam JJ, Cuvillon P, Capdevila X et al. Postoperative analgesia by femoral nerve block with ropivacaine 0.2% after major knee surgery: continuous versus patient-controlled techniques. *Reg Anesth Pain Med* 2002;27:604-11.
32. Badner NH, Bourne RB, Rorabeck CH et al. Intra-articular injection of bupivacaine in knee-replacement operations. Results of use for analgesia and for preemptive blockade. *J Bone Joint Surg Am* 1996;78:734-8.
33. Browne C, Copp S, Reden L, et al. Bupivacaine bolus injection versus placebo for pain management following total knee arthroplasty. *J Arthroplasty* 2004;19:377-80.
34. Mauerhan DR, Campbell M, Miller JS, et al. Intra-articular morphine and/or bupivacaine in the management of pain after total knee arthroplasty. *J Arthroplasty* 1997;12:546-52.
35. Ritter MA, Koehler M, Keating EM, et al. Intra-articular morphine and/or bupivacaine after total knee replacement. *J Bone Joint Surg Br* 1999;81:301-3.
36. Tanaka N, Sakahashi H, Sato E et al. The efficacy of intra-articular analgesia after total knee arthroplasty in patients with rheumatoid arthritis and in patients with osteoarthritis. *J Arthroplasty* 2001;16:306-11.
37. Isaac D, Falode T, Liu P et al. Accelerated rehabilitation after total knee replacement. *Knee* 2005;12:346-50.
38. Bianconi M, Ferraro L, Traina GC et al. Pharmacokinetics and efficacy of ropivacaine continuous wound instillation after joint replacement surgery. *Br J Anaesth* 2003;91:830-5.
39. Nechleba J, Rogers V, Cortina G et al. Continuous intra-articular infusion of bupivacaine for postoperative pain following total knee arthroplasty. *J Knee Surg* 2005;18:197-202.
40. Pulido PA, Colwell J, Hoenecke J et al. The efficacy of continuous bupivacaine infiltration for pain management following orthopaedic knee surgery: anterior cruciate ligament reconstruction and total knee arthroplasty. *Orthop Nurs* 2002;21:31-6.

> AKADEMISKE AFHANDLINGER

Afdelingslæge Flemming Møller:

Quantification and characterization of fixational eye movements in normal test persons and in patients with macular disease

Disputats

Forf.s adresse: Rosenvænget 4, DK-5250 Odense SV.

E-mail: f.moeller@dadlnet.dk

Forsvaret finder sted den 9. juni 2006, kl. 13.00, Auditorium U110, Syddansk Universitet, Campusvej 55, Odense M.

Opponenter: Monica Lövestam-Adrian, Sverige, Erik Krogh og Søren Sindrup.

Cand.scient. Marianne Jensby Nielsen:

Molekylær karakterisering af haptoglobin-hæmoglobin-komplekset og dets receptor

Ph.d.-afhandling

Forf.s adresse: Institut for Medicinsk Biokemi, Ole Worms Allé, Bygning 1170, DK-8000 Århus C.

E-mail: mjn@biokemi.au.dk

Forsvaret finder sted den 19. maj 2006, kl. 14.00, Auditorium 6, Institut for Medicinsk Biokemi, Ole Worms Allé, Bygning 1170, Aarhus Universitet.

Bedømmere: Bente Vilse, Lars Bo Nielsen og Uffe Holmskov.

Vejleder: Søren K. Moestrup.

Find referater af denne uges akademiske afhandlinger på
www.ugeskriftet.dk under

>Seneste nummer >Akademiske afhandlinger